

SINGULAR SUPPORTS FOR CHARACTER SHEAVES ON A GROUP COMPACTIFICATION

XUHUA HE AND GEORGE LUSZTIG

ABSTRACT. Let G be a semisimple adjoint group over $\mathbf{C}$ and $\bar{G}$ be the De Concini-Procesi completion of G . In this paper, we define a Lagrangian subvariety Λ of the cotangent bundle of $\bar{G}$ such that the singular support of any character sheaf on $\bar{G}$ is contained in Λ .

1.1. In this paper all algebraic varieties are assumed to be over a fixed algebraically closed field of characteristic 0.

If X is a smooth variety, let T^*X be the cotangent bundle of X . For any morphism $\alpha : X \rightarrow Y$ of smooth varieties and $x \in X$, we write $\alpha^* : T_{\alpha(x)}^*Y \rightarrow T_x^*X$ for the map induced by α . If moreover, $\alpha : X \rightarrow Y$ is a locally trivial fibration with smooth connected fibres and Λ is a closed Lagrangian subvariety of T^*Y , then let $\alpha^\star(\Lambda) = \cup_{x \in X} \alpha^*(\Lambda \cap T_{\alpha(x)}^*Y) \subset T^*X$. Then

(a) $\alpha^\star(\Lambda)$ is a closed Lagrangian subvariety of T^*X . Moreover, the set of irreducible components of Λ is naturally in bijection with the set of irreducible components of $\alpha^\star(\Lambda)$.

Let X, Y be smooth irreducible varieties and let $\alpha : X \rightarrow Y$ be a principal P -bundle for a free action of a connected linear algebraic group P on X .

(b) If Λ' is a closed Lagrangian subvariety of T^*X stable under the P -action then $\Lambda' = \alpha^\star(\Lambda)$ for a unique closed Lagrangian subvariety Λ of T^*Y .

Let X be a smooth irreducible variety and let $i : Y \rightarrow X$ be the inclusion of a locally closed smooth irreducible subvariety. Let Λ be a closed Lagrangian subvariety of T^*Y . Let $i_\star(\Lambda)$ be the subset of T^*X consisting of all $\xi \in T_x^*X$ such that $x \in Y$ and the image of ξ under the obvious surjective map $T_x^*X \rightarrow T_x^*Y$ belongs to $\Lambda \cap T_x^*Y$. Note that

(c) $i_\star(\Lambda)$ is a locally closed Lagrangian subvariety of T^*X . Moreover, the set of irreducible components of $i_\star(\Lambda)$ is naturally in bijection with the set of irreducible components of Λ .

2000 *Mathematics Subject Classification.* 20G99.

X. H. is supported by NSF grant DMS-0111298. G.L. is supported in part by NSF grant DMS-0243345.

For an algebraic variety X we write $\mathcal{D}(X)$ for the bounded derived category of constructible $\bar{\mathbf{Q}}_l$ -sheaves on X where l is a fixed prime number. For X smooth and $C \in \mathcal{D}(X)$, we denote by $SS(X)$ the singular support of C (a closed Lagrangian subvariety of T^*X). Let A be a connected linear algebraic group acting on a smooth variety X and let B be a connected subgroup of A . Let $\mu_A : T^*X \rightarrow \text{Lie}(A)^*$ be the moment map of the A -action on X . Consider the diagram $X \xleftarrow{pr_1} A \times X \xrightarrow{pr_2} A \times_B X \xrightarrow{\pi} X$ where B acts on $A \times X$ by $b : (a, x) \mapsto (ab^{-1}, bx)$, $A \times_B X$ is the quotient space and $\pi(a, x) = ax$. Then for any B -equivariant perverse sheaf C on X there is a well defined perverse sheaf C' on $A \times_B X$ such that $pr_2^*C' = pr_1^*C$ up to a shift. We set $\Gamma_B^A(C) = \pi_*C' \in \mathcal{D}(X)$. By [MV, 1.2] we have

$$(d) \ SS(\Gamma_B^A(C)) \subset \overline{A \cdot SS(C)}.$$

On the other hand, we have

$$(e) \ SS(C) \subset \mu_B^{-1}(0).$$

Indeed, if $p_1 : B \times X \rightarrow X$ is the action and $p_2 : B \times X \rightarrow X$ is the second projection we have $p_1^*(C) = p_2^*(C)$. Hence $SS(p_1^*(C)) = SS(p_2^*(C))$. Using [KS, 4.1.2] we can rewrite this as $p_1^\star(SS(C)) = p_2^\star(SS(C))$. Hence if $x \in X$ and $\xi \in T_x^*X \cap SS(C)$ then the image of ξ under the map $T_x^*X \rightarrow T_1^*(B)$ induced by $B \rightarrow X, b \mapsto bx$ is 0. This proves (e).

1.2. Let G be a connected reductive algebraic group. Let $\mathfrak{g} = \text{Lie}(G)$. Let $\mathcal{N}$ be the variety of nilpotent elements in $\mathfrak{g}^*$. Let B be a Borel subgroup of G . Let K be a closed connected subgroup of G . Then $B_K = B \cap K$ is a parabolic subgroup of K . Assume that G acts on a smooth variety X . Let C be a B_K -equivariant perverse sheaf on X ; assume also that there exists a finite covering $a : \tilde{B} \rightarrow B$ such that C is $\tilde{B}$ -equivariant for the $\tilde{B}$ -action $\tilde{b} : x \mapsto a(\tilde{b})x$ on X . By 1.1(e) we have $\mu_{\tilde{B}}(SS(C)) = 0$. Since $\text{Lie}(\tilde{B}) = \text{Lie}(B)$ we then have $\mu_B(SS(C)) = 0$. It follows that $\mu_G(SS(C))$ is contained in the kernel of the obvious map $\mathfrak{g}^* \rightarrow \text{Lie}(B)^*$ hence is contained in $\mathcal{N}$. Since $\mathcal{N}$ is stable under the coadjoint action we have $\mu_G(K \cdot SS(C)) = K\mu_G(SS(C)) \subset \mathcal{N}$. Using this together with 1.1(d) and the fact that $\mu_G^{-1}(\mathcal{N})$ is closed in T^*X we see that $SS(\Gamma_{B_K}^K(C)) \subset \mu_G^{-1}(\mathcal{N})$. Applying 1.1(e) to $\Gamma_{B_K}^K(C)$, K instead of C, B we see that $SS(\Gamma_{B_K}^K(C)) \subset \mu_K^{-1}(0) = \mu_G^{-1}(\text{Lie}(K)^\perp)$ where $\text{Lie}(K)^\perp \subset \mathfrak{g}^*$ is the annihilator of $\text{Lie}(K) \subset \mathfrak{g}$. Thus we have

$$(a) \quad SS(\Gamma_{B_K}^K(C)) \subset \mu_G^{-1}(\text{Lie}(K)^\perp \cap \mathcal{N}).$$

1.3. We now replace $G, \mathfrak{g}, B, K, B_K, X, C$ by $G \times G, \mathfrak{g} \times \mathfrak{g}, B \times B, G_\Delta, B_\Delta, X', C'$ where $G_\Delta = \{(g, g') \in G \times G; g = g'\}$, $B_\Delta = \{(g, g') \in B \times B; g = g'\}$, X' is a smooth variety with a given action of $G \times G$ and C' is a B_Δ -equivariant perverse sheaf on X' ; we assume that there exists a finite covering $a' : \tilde{B}' \rightarrow B \times B$ such that C' is $\tilde{B}'$ -equivariant for the $\tilde{B}'$ -action $\tilde{b}' : x' \mapsto a'(\tilde{b}')x'$ on X' . We have the following special case of 1.2(a):

$$(a) \quad SS(\Gamma_{B_\Delta}^{G_\Delta}(C')) \subset \mu_{G \times G}^{-1}(\mathcal{N}^-)$$

where

$$\mathcal{N}^- = \{(f, f') \in \mathfrak{g}^* \times \mathfrak{g}^*; f + f' = 0, f, f' \text{ nilpotent} \}.$$

1.4. Let $\mathbf{W}$ be the Weyl group of G and let $\mathbf{I}$ be the set of simple reflections in $\mathbf{W}$. Let G' be a possibly disconnected algebraic group with identity component G and with a given connected component D . Now $G \times G$ acts transitively on D by $(g_1, g_2) : g \mapsto g_1 g g_2^{-1}$. Hence the moment map $\mu_{G \times G} : T^*D \rightarrow \mathfrak{g}^* \times \mathfrak{g}^*$ is well defined. In [L1, 4.5] a class of perverse sheaves (called character sheaves) on D is introduced. These appear as constituents of some perverse cohomology sheaf of $\Gamma_{B\Delta}^{G\Delta}(C')$ for some C' as in 1.3 (with $X' = D$). Hence from 1.3(a) we deduce:

(a) *If K is a parabolic character sheaf on D then $SS(K) \subset \mu_{G \times G}^{-1}(\mathcal{N}^-)$.*

In the case where $G' = G = D$ a statement close to (a) appears in [MV, 2.8] (where it is attributed to the second author) and in [Gi].

1.5. We preserve the setup of 1.4. For any $J \subset \mathbf{I}$ let $\mathcal{P}_J$ be the set of parabolic subgroups of G of type J . In particular $\mathcal{P}_\emptyset$ is the set of Borel subgroups of G . For $J \subset \mathbf{I}$ let $\mathbf{W}_J$ be the subgroup of $\mathbf{W}$ generated by J ; let $\mathbf{W}^J$ (resp. ${}^J\mathbf{W}$) be the set of all $w \in \mathbf{W}$ such that w has minimal length among the elements in $\mathbf{W}_J w$ (resp. $w\mathbf{W}_J$). Let $\delta : \mathbf{W} \xrightarrow{\sim} \mathbf{W}$ be the isomorphism such that $\delta(\mathbf{I}) = \mathbf{I}$ and such that $J \subset \mathbf{I}, P \in \mathcal{P}_J, g \in D \implies gPg^{-1} \in \mathcal{P}_{\delta(J)}$. Following [L2, 8.18], for $J, J' \subset \mathbf{I}$ and $y \in {}^{J'}\mathbf{W} \cap \mathbf{W}^J$ such that $\text{Ad}(y)(\delta(J)) = J'$ we set

$$Z_{J,y,\delta} = \{(P, P', gU_P); P \in \mathcal{P}_J, P' \in \mathcal{P}_{J'}, g \in D, \text{pos}(P', gPg^{-1}) = y\}.$$

Now $G \times G$ acts (transitively) on $Z_{J,y,\delta}$ by

$$(g_1, g_2) : (P, P', gU_P) \mapsto (g_2 P g_2^{-1}, g_1 P' g_1^{-1}, g_1 g g_2^{-1}).$$

Hence the moment map $\mu_{G \times G} : T^*Z_{J,y,\delta} \rightarrow \mathfrak{g}^* \times \mathfrak{g}^*$ is well defined. In [L2, Section 11] a class of perverse sheaves (called parabolic character sheaves) on $Z_{J,y,\delta}$ is introduced. These appear as constituents of some perverse cohomology sheaf of $\Gamma_{B\Delta}^{G\Delta}(C')$ for some C' as in 1.3 (with $X' = Z_{J,y,\delta}$). Hence from 1.3(a) we deduce:

(a) *If K is a parabolic character sheaf on $Z_{J,y,\delta}$ then $SS(K) \subset \mu_{G \times G}^{-1}(\mathcal{N}^-)$.*

When $J = \mathbf{I}$, this reduces to 1.4(a).

1.6. Assume that G is adjoint. Let $\bar{G}$ be the De Concini-Procesi compactification of G . Then $G \times G$ acts naturally on $\bar{G}$ extending continuously the action $(g_1, g_2) : g \mapsto g_1 g g_2^{-1}$ of $G \times G$ on G . Hence the moment map $\mu_{G \times G} : T^*\bar{G} \rightarrow \mathfrak{g}^* \times \mathfrak{g}^*$ is well defined. In [L2] a class of perverse sheaves (called parabolic character sheaves) on $\bar{G}$ is introduced. It has been shown by He [H2] and by Springer (unpublished) that any parabolic character sheaf on $\bar{G}$ appears as a constituent of some perverse cohomology sheaf of $\Gamma_{B\Delta}^{G\Delta}(C')$ for some C' as in 1.3 (with $X' = \bar{G}$). Hence from 1.3(a) we deduce:

(a) *If K is a parabolic character sheaf on $\bar{G}$ then $SS(K) \subset \mu_{G \times G}^{-1}(\mathcal{N}^-)$.*

1.7. In the setup of 1.4 let $\Lambda(D) = \mu_{G \times G}^{-1}(\mathcal{N}^-)$. We want to describe the variety $\Lambda(D)$. For $g \in D$ let I_g be the isotropy group at g of the $G \times G$ -action on D that is, $I_g = \{(g_1, g_2) \in G \times G; g_2 = g^{-1}g_1g\}$. We have $\text{Lie}(I_g) = \{(y_1, y_2) \in \mathfrak{g} \times \mathfrak{g}; y_2 = \text{Ad}(g)^{-1}(y_1)\}$ and the annihilator of $\text{Lie}(I_g)$ in $\mathfrak{g}^* \times \mathfrak{g}^*$ is $\text{Lie}(I_g)^\perp = \{(z_1, z_2) \in \mathfrak{g}^* \times \mathfrak{g}^*; z_1 + \text{Ad}(g)(z_2) = 0\}$. This may be identified with the fibre of T^*D at g . Then

$$\begin{aligned} \Lambda(D) &= \{(g, z_1, z_2) \in D \times \mathfrak{g}^* \times \mathfrak{g}^*; z_1 + \text{Ad}(g)(z_2) = 0, z_1 + z_2 = 0, z_2 \in \mathcal{N}\} \\ &= \{(g, z, -z); g \in D, z \in \mathcal{N}, \text{Ad}(g)(z) = z\} = \sqcup_{\mathcal{O}} X_{\mathcal{O}} \end{aligned}$$

where $\mathcal{O}$ runs over the (finite) set of $\text{Ad}(G)$ -orbits on $\mathcal{N}$ which are normalized by some element of D and $X_{\mathcal{O}} = \{(g, z, -z); g \in D, z \in \mathcal{O}, \text{Ad}(g)(z) = z\}$. We pick $\xi \in \mathcal{O}$ and let $\mathcal{Z}' = \{h \in G'; \text{Ad}(h)\xi = \xi\}$, $\mathcal{Z} = \{h \in G; \text{Ad}(h)\xi = \xi\}$. Let $\underline{\mathcal{Z}}'$ (resp. $\underline{\mathcal{Z}}$) be the group of connected components of $\mathcal{Z}'$ (resp. $\mathcal{Z}$). Then $\underline{\mathcal{Z}}'$ is a finite group and $\underline{\mathcal{Z}}$ is a subgroup of $\underline{\mathcal{Z}}'$. Let $\underline{\mathcal{Z}}_1$ be the set of connected components of $\mathcal{Z}'$ that are contained in D . Then $\underline{\mathcal{Z}}_1$ is a subset of $\mathcal{Z}'$; also, $\underline{\mathcal{Z}}$ acts on $\underline{\mathcal{Z}}_1$ by conjugation inside $\underline{\mathcal{Z}}'$. Let $F_{\mathcal{O}}^D$ be the set of orbits of this action. Note that $F_{\mathcal{O}}^D$ is independent (up to unique isomorphism) of the choice of ξ .

Let $\tilde{X} = \{(g, r) \in D \times G; r^{-1}gr \in \mathcal{Z}'\}$. Then $\mathcal{Z}$ acts freely on $\tilde{X}$ by $h : (g, r) \mapsto (g, rh^{-1})$ and we have an isomorphism $\mathcal{Z} \backslash \tilde{X} \xrightarrow{\sim} X_{\mathcal{O}}$, $(g, r) \mapsto (g, \text{Ad}(r)\xi)$. By the change of variable $(g, r) \mapsto (g', r)$, $g' = r^{-1}gr$, $\tilde{X}$ becomes $\{(g', r); g' \in \mathcal{Z}' \cap D, r \in G\}$. In the new coordinates, the free action of $\mathcal{Z}$ on $\tilde{X}$ is $h : (g', r) \mapsto (hg'h^{-1}, rh^{-1})$. We see that $\tilde{X}$ is smooth of pure dimension $\dim(\mathcal{Z} \times G)$ and its connected components are indexed naturally by $\underline{\mathcal{Z}}_1$ (the connected component containing (g', r) is indexed by the image of g' in $\underline{\mathcal{Z}}_1$). The action of $\mathcal{Z}$ on $\tilde{X}$ permutes the connected components of $\tilde{X}$ according to the action of $\underline{\mathcal{Z}}$ on $\underline{\mathcal{Z}}_1$ considered above. We see that $X_{\mathcal{O}} = \mathcal{Z} \backslash \tilde{X}$ is smooth of pure dimension $\dim G$ and its connected components are indexed naturally by the set $F_{\mathcal{O}}^D$.

We see that $\Lambda(D)$ can be partitioned into finitely many locally closed, irreducible, smooth subvarieties of dimension $\dim G$, indexed by the finite set $F(D) := \sqcup_{\mathcal{O}} F_{\mathcal{O}}^D$. In particular, $\Lambda(D)$ has pure dimension $\dim G$. More precisely, one checks that

(a) $\Lambda(D)$ is a closed Lagrangian subvariety of T^*D .

1.8. In the setup of 1.5 we set $\Lambda(Z_{J,y,\delta}) = \mu_{G \times G}^{-1}(\mathcal{N}^-)$. We want to describe the variety $\Lambda(Z_{J,y,\delta})$.

Following [L2, 8.18] we consider the partition $Z_{J,y,\delta} = \sqcup_{\mathbf{s}} Z_{J,y,\delta}^{\mathbf{s}}$ where $Z_{J,y,\delta}^{\mathbf{s}}$ are certain locally closed smooth irreducible G_{Δ} -stable subvarieties of $Z_{J,Y,\delta}$ indexed by the elements $\mathbf{s}$ of a finite set $S(J, \text{Ad}(y)\delta)$ which is in canonical bijection with $J' \mathbf{W}$, see [L1, 2.5]. Note that each $\mathbf{s}$ is a sequence $(J_n, J'_n, u_n)_{n \geq 0}$ where J_n, J'_n are subsets of $\mathbf{I}$ such that J_n, J'_n are independent of n for large n and $u_n \in \mathbf{W}$ is 1 for large n .

We wish to define a Lagrangian subvariety $\Lambda(Z_{J,y,\delta}^{\mathbf{s}})$ of $T^*(Z_{J,y,\delta}^{\mathbf{s}})$.

Assume first that $\mathbf{s}$ is such that $J_n = J, J'_n = J', u_n = 1$ for all n . In this case we have $J = J'$. Let $P \in \mathcal{P}_J$ and let L be a Levi subgroup of P . Then $\mathbf{d}_{\mathbf{s}} = \{g \in D; gLg^{-1} = L, \text{pos}(P, gPg^{-1}) = y\}$ is a connected component of the algebraic group $N_{G'}(L)$ with identity component L . Hence $\Lambda(\mathbf{d}_{\mathbf{s}}) \subset T^*\mathbf{d}_{\mathbf{s}}$ is defined as in 1.7. We have a diagram $Z_{J,y,\delta}^{\mathbf{s}} \xleftarrow{\alpha} G \times \mathbf{d}_{\mathbf{s}} \xrightarrow{pr_2} \mathbf{d}_{\mathbf{s}}$ where $\alpha(h, g) = (hPh^{-1}, hPh^{-1}, U_{hPh^{-1}}hgh^{-1}U_{hPh^{-1}})$. Note that α is a principal P -bundle where P acts on $G \times \mathbf{d}_{\mathbf{s}}$ by $p : (h, g) = (hp^{-1}, \bar{p}g\bar{p}^{-1})$ (we denote the canonical homomorphism $P \rightarrow L$ by $\bar{p}$). Let $\Lambda' = pr_2^*\Lambda(\mathbf{d}_{\mathbf{s}}) \subset T^*(G \times \mathbf{d}_{\mathbf{s}})$. By 1.1(a) and 1.7(a), Λ' is a closed Lagrangian subvariety of $T^*(G \times \mathbf{d}_{\mathbf{s}})$. It is clearly stable under the natural action of P on $T^*(G \times \mathbf{d}_{\mathbf{s}})$ (since $\Lambda(\mathbf{d}_{\mathbf{s}})$ is L -stable). By 1.1(b) there is a unique Lagrangian subvariety Λ'' of $T^*(Z_{J,y,\delta}^{\mathbf{s}})$ such that $\alpha^{\star}\Lambda'' = \Lambda'$. We set $\Lambda(Z_{J,y,\delta}^{\mathbf{s}}) = \Lambda''$.

We now consider a general $\mathbf{s} = (J_n, J'_n, u_n)_{n \in \mathbf{N}}$. For any $r \in \mathbf{N}$ let $\mathbf{s}_r = (J_n, J'_n, u_n)_{n \geq r}$, $y_r = u_{r-1}^{-1} \dots u_1^{-1} u_0^{-1} y$. Then $Z_{J_r, y_r, \delta}^{\mathbf{s}_r}$ is defined and we have a canonical map $f_r : Z_{J,y,\delta}^{\mathbf{s}} \rightarrow Z_{J_r, y_r, \delta}^{\mathbf{s}_r}$ (a composition of affine space bundles, see [L2, 8.20(a)]). Moreover for sufficiently large r , $\mathbf{s}_r, J_r, y_r, f_r$ are independent of r ; we write $\mathbf{s}_{\infty}, J_{\infty}, y_{\infty}, f_{\infty}$ instead of $\mathbf{s}_r, J_r, y_r, f_r$. Note also that $\mathbf{s}_{\infty}, J_{\infty}, y_{\infty}$ are of the type considered earlier, so that $\Lambda(Z_{J_{\infty}, y_{\infty}, \delta}^{\mathbf{s}_{\infty}})$ is defined as above. We set $\Lambda(Z_{J,y,\delta}^{\mathbf{s}}) = f_{\infty}^{\star}(\Lambda(Z_{J_{\infty}, y_{\infty}, \delta}^{\mathbf{s}_{\infty}}))$.

We now define

$$\Lambda'(Z_{J,y,\delta}) = \sqcup_{\mathbf{s} \in S(J, \text{Ad}(y)\delta)} (i_{\mathbf{s}})^{\star}(\Lambda(Z_{J,y,\delta}^{\mathbf{s}}))$$

where $i_{\mathbf{s}} : Z_{J,y,\delta}^{\mathbf{s}} \rightarrow Z_{J,y,\delta}$ is the inclusion. From 1.1(c) we see that $\Lambda'(Z_{J,y,\delta})$ is a finite union of locally closed Lagrangian subvarieties of $T^*(Z_{J,y,\delta})$.

We state the following result:

Proposition 1.9. *We have $\Lambda(Z_{J,y,\delta}) = \Lambda'(Z_{J,y,\delta})$. In particular, $\Lambda'(Z_{J,y,\delta})$ is closed in $T^*(Z_{J,y,\delta})$ and $\Lambda(Z_{J,y,\delta})$ is a Lagrangian subvariety of $T^*(Z_{J,y,\delta})$.*

Let $x \in Z_{J,y,\delta}^{\mathbf{s}}$. We will show that $\Lambda(Z_{J,y,\delta}) \cap T_x^*(Z_{J,y,\delta}) = \Lambda'(Z_{J,y,\delta}) \cap T_x^*(Z_{J,y,\delta})$.

We identify $\mathfrak{g}$ with $\mathfrak{g}^*$ via a G -invariant symmetric bilinear form. Choose an element g_0 in D that normalizes B and a maximal torus T of B . Let P_J be the unique element in $\mathcal{P}_J$ that contains B . For $w \in W$, we choose a representative $\dot{w}$ of w in $N(T)$. Set $h_{J,y,\delta} = (P_J, \dot{y}^{-1} P_{J'}, U_{\dot{y}^{-1} P_{J'}} g_0 U_{P_J}) \in Z_{J,y,\delta}$. The element $\mathbf{s} \in S(J, \text{Ad}(y)\delta)$ corresponds to an element $w' \in {}^{J'}W$ under the bijection in [L1, 2.5]. Set $w = (w')^{-1} y \in w \in W^{\delta(J)}$. By [H1, 1.10],

$$Z_{J,y,\delta}^{\mathbf{s}} = G_{\Delta}(P_{J_{\infty}} \dot{w}, 1) \cdot h_{J,y,\delta} = G_{\Delta}(L_{J_{\infty}} \dot{w}, 1) \cdot h_{J,y,\delta}.$$

Note that $\Lambda(Z_{J,y,\delta})$ and $\Lambda'(Z_{J,y,\delta})$ are stable under the action of G_{Δ} . Then we may assume that $x = (l_1 \dot{w}, 1) \cdot h_{J,y,\delta}$ for some $l_1 \in L_{J_{\infty}}$. By [H1, 1.10(2)], $(P_{J_{\infty}})_{\Delta} \cdot (L_{J_{\infty}}, 1)x = (P_{J_{\infty}}, 1) \cdot x$, where $(P_{J_{\infty}})_{\Delta} = \{(g, g') \in P_{J_{\infty}} \times P_{J_{\infty}}; g = g'\}$. For $u \in U_{P_{J_{\infty}}}$, $(u, 1) \cdot x = (u'l, u') \cdot x$ for some $u' \in U_{P_{J_{\infty}}}$ and $l \in L_{J_{\infty}}$. Note that

$(U_{P_{J_\infty}}, U_{P_{J_\infty}}) \cdot (l, 1) \cdot x = (U_{P_{J_\infty}} l, 1) \cdot x$ and $(P_{J_\infty}, 1) \cdot x \cong U_{P_{J_\infty}} \times (L_{J_\infty}, 1) \cdot x$. Thus $(l, 1) \cdot x = x$ and $(U_{P_{J_\infty}}, 1) \cdot x \subset (P_{J_\infty})_\Delta \cdot x$. Therefore, for $(f, -f) \in \mu_{G \times G}(T_x^*(Z_{J,y,\delta}))$, we have that $(f, -f)(\text{Lie}(U_{P_{J_\infty}}), 0) = (f, -f)(\text{Lie}(P_{J_\infty})_\Delta) = 0$, i. e., $f \in \text{Lie}(P_{J_\infty})$. In particular, f is nilpotent if and only if the image of f under $\text{Lie}(P_{J_\infty}) \rightarrow \text{Lie}(P_{J_\infty})/\text{Lie}(U_{P_{J_\infty}}) \cong \text{Lie}(L_{J_\infty})$ is nilpotent.

Hence $\mu_{G \times G}(\Lambda(Z_{J,y,\delta}) \cap T_x^*(Z_{J,y,\delta}^s))$ consists elements of the form $(u+l, -u-l)$ with $u \in \text{Lie}(U_{P_{J_\infty}})$, l nilpotent in $\text{Lie}(L_{J_\infty})$ and $(u, -u)I_x = (l, -l)I_x = 0$, where I_x is the Lie subalgebra of the isotropic subgroup of $G \times G$ at point x .

Denote by N_x the stalk at point x of the conormal bundle $N_{Z_{J,y,\delta}^s}^*(Z_{J,y,\delta})$. Since $Z_{J,y,\delta}^s = G_\Delta(P_{J_\infty}, 1)h_{J,y,\delta}$, we have

$$\mu_{G \times G}(N_x) = \{(u, -u); u \in \text{Lie}(U_{P_{J_\infty}}), (u, -u)I_x = 0\}.$$

Let $p_x : T_x^*(Z_{J,y,\delta}) \rightarrow T_x^*(Z_{J,y,\delta}^s) \cong T_x^*(Z_{J,y,\delta})/N_x$ be the obvious surjective map. Then $\Lambda(Z_{J,y,\delta}) \cap T_x^*(Z_{J,y,\delta}) = p_x^{-1}(p_x(\Lambda(Z_{J,y,\delta}) \cap T_x^*(Z_{J,y,\delta})))$. Note that

$$I_x = \{(u_1 + \text{Ad}(l_1 \dot{w} g_0)l, u_2 + l); u_1 \in \text{Ad}(l_1 \dot{w} g_0^{-1})\text{Lie}(U_{P_{J'}}), u_2 \in \text{Lie}(U_{P_J}), l \in \text{Lie}(L_J)\}. \blacksquare$$

Thus for $l \in \text{Lie}(L_{J_\infty})$, $(l, -l)I_x = 0$ if and only if $\text{Ad}(l_1 \dot{w} g_0)l = l$.

We identify $T_x^*(Z_{J,y,\delta})$ with $\mu_{G \times G}(T_x^*(Z_{J,y,\delta})) \subset \mathfrak{g} \times \mathfrak{g}$ and regard $T_x^*(Z_{J,y,\delta}^s)$ as a subspace of $(\mathfrak{g} \times \mathfrak{g})/N_x$. Set $M = \{(l, -l); l \in \text{Lie}(L_{J_\infty}), \text{Ad}(l_1 \dot{w} g_0)l = l\} \subset \mathfrak{g} \times \mathfrak{g}$. Then

$$(1) \ p_x(\Lambda(Z_{J,y,\delta}) \cap T_x^*(Z_{J,y,\delta})) = (M + N_x)/N_x.$$

Now consider the commuting diagram

$$\begin{array}{ccc} (P_{J_\infty})_\Delta(L_{J_\infty} \dot{w}, 1) \cdot h_{J,y,\delta} & \xrightarrow{i_1} & Z_{J,y,\delta}^s \\ f'_\infty \downarrow & & f_\infty \downarrow \\ (P_{J_\infty})_\Delta(L_{J_\infty} \dot{w}, 1) \cdot h_{J_\infty, y_\infty, \delta} & \xrightarrow{i_2} & Z_{J_\infty, y_\infty, \delta}^s \end{array}$$

where i_1, i_2 are inclusions and f'_∞ is the restriction of f_∞ . Let $x' = f_\infty(x) \in Z_{J_\infty, y_\infty, \delta}^s$. Since f'_∞ is $P_{J_\infty} \times P_{J_\infty}$ -invariant, we have the following commuting diagram

$$\begin{array}{ccc} T_{x'}^*((P_{J_\infty})_\Delta(L_{J_\infty} \dot{w}, 1) \cdot h_{J_\infty, y_\infty, \delta}) & \xrightarrow{{}^1\mu_{P_{J_\infty} \times P_{J_\infty}}} & \text{Lie}(P_{J_\infty})^* \times \text{Lie}(P_{J_\infty})^* \\ (f'_\infty)^* \downarrow & & id \downarrow \\ T_x^*((P_{J_\infty})_\Delta(L_{J_\infty} \dot{w}, 1) \cdot h_{J,y,\delta}) & \xrightarrow{{}^2\mu_{P_{J_\infty} \times P_{J_\infty}}} & \text{Lie}(P_{J_\infty})^* \times \text{Lie}(P_{J_\infty})^*, \end{array}$$

where ${}^1\mu_{P_{J_\infty} \times P_{J_\infty}}$ and ${}^2\mu_{P_{J_\infty} \times P_{J_\infty}}$ are the moment maps. Since the actions of $P_{J_\infty} \times P_{J_\infty}$ on $(P_{J_\infty})_\Delta(L_{J_\infty} \dot{w}, 1) \cdot h_{J,y,\delta}$ and $(P_{J_\infty})_\Delta(L_{J_\infty} \dot{w}, 1) \cdot h_{J_\infty, y_\infty, \delta}$ are transitive, ${}^1\mu_{P_{J_\infty} \times P_{J_\infty}}$ and ${}^2\mu_{P_{J_\infty} \times P_{J_\infty}}$ are injective.

Set $\Lambda_{x'}(Z_{J_\infty, y_\infty, \delta}^{\mathbf{s}_\infty}) = \Lambda(Z_{J_\infty, y_\infty, \delta}^{\mathbf{s}_\infty}) \cap T_{x'}^*(Z_{J_\infty, y_\infty, \delta}^{\mathbf{s}_\infty})$. Then

$${}^2\mu_{P_{J_\infty} \times P_{J_\infty}}(i_2^*(\Lambda_{x'}(Z_{J_\infty, y_\infty, \delta}^{\mathbf{s}_\infty}))) = {}^1\mu_{P_{J_\infty} \times P_{J_\infty}}((f'_\infty)^* i_2^*(\Lambda_{x'}(Z_{J_\infty, y_\infty, \delta}^{\mathbf{s}_\infty}))) = M.$$

Here we identify $\text{Lie}(P_{J_\infty})^*$ with $\text{Lie}(P_{J_\infty}^-)$ via the symmetric bilinear form. Moreover, $i_1^* f_\infty^* = (f'_\infty)^* i_2^*$ maps $\Lambda_{x'}(Z_{J_\infty, y_\infty, \delta}^{\mathbf{s}_\infty})$ bijectively onto its image. Therefore ${}^1\mu_{P_{J_\infty} \times P_{J_\infty}}(i_1^*(\Lambda(Z_{J, y, \delta}^{\mathbf{s}}) \cap T_x^*(Z_{J, y, \delta}^{\mathbf{s}}))) = M$ and i_1^* maps $\Lambda(Z_{J, y, \delta}^{\mathbf{s}}) \cap T_x^*(Z_{J, y, \delta}^{\mathbf{s}})$ bijectively onto its image. In other words,

$$(2) \Lambda(Z_{J, y, \delta}^{\mathbf{s}}) \cap T_x^*(Z_{J, y, \delta}^{\mathbf{s}}) = (M + N_x)/N_x.$$

Combining (1) and (2), $p_x(\Lambda(Z_{J, y, \delta}) \cap T_x^*(Z_{J, y, \delta})) = \Lambda(Z_{J, y, \delta}^{\mathbf{s}}) \cap T_x^*(Z_{J, y, \delta}^{\mathbf{s}})$. The proposition is proved.

Corollary 1.10. *The set of irreducible components of $\Lambda(Z_{J, y, \delta})$ is in natural bijection with $\sqcup_{\mathbf{s} \in S(J, \text{Ad}(y)\delta)} F(\mathbf{d}_{\mathbf{s}_\infty})$ (notation of 1.7, 1.8).*

1.11. In the setup of 1.6 let $\Lambda(\bar{G}) = \mu_{G \times G}^{-1}(\mathcal{N}^-)$. We want to describe the variety $\Lambda(Z_{J, y, \delta})$. As in [L2, 12.3], we have $\bar{G} = \sqcup_{J \subset \mathbf{I}} G_J$ where G_J are the various $G \times G$ -orbits in $\bar{G}$; moreover we may identify $G_J = T_J \backslash Z_{J, y_J, 1}$ where y_J is the longest element in $\mathbf{W}^J$ and T_J is a torus acting freely on $Z_{J, y_J, 1}$. Let $a_J : Z_{J, y_J, 1} \rightarrow G_J$ be the canonical map.

For each J let $\mu_{G \times G; J} : G_J \rightarrow \mathfrak{g}^* \times \mathfrak{g}^*$ be the moment map of the restriction of the $G \times G$ -action on $\bar{G}$ to G_J . Let $i_J : G_J \rightarrow \bar{G}$ be the inclusion. From the definitions, we have

$$(a) \Lambda(\bar{G}) = \sqcup_{J \subset \mathbf{I}} (i_J)_\star \mu_{G \times G; J}^{-1}(\mathcal{N}^-)$$

$$\text{and } \Lambda(Z_{J, y_J, 1}) = a_J^\star(\mu_{G \times G; J}^{-1}(\mathcal{N}^-)).$$

Since $\Lambda(Z_{J, y_J, 1})$ is a T_J -stable Lagrangian subvariety of $T^*(Z_{J, y_J, 1})$ (see 1.9), it follows that $\mu_{G \times G; J}^{-1}(\mathcal{N}^-)$ is a Lagrangian subvariety of $T^*(G_J)$. Hence, using 1.1(c), we see that $(i_J)_\star \mu_{G \times G; J}^{-1}(\mathcal{N}^-)$ is a Lagrangian subvariety of $T^*\bar{G}$. Using this and (a) we see that

$$(b) \Lambda(\bar{G}) \text{ is a Lagrangian subvariety of } T^*\bar{G}.$$

From the previous proof we see that

$$(c) \text{ The set of irreducible components of } \Lambda(\bar{G}) \text{ is in natural bijection with } \sqcup_{J \subset \mathbf{I}} \sqcup_{\mathbf{s} \in S(J, \text{Ad}(y_J))} F(\mathbf{d}_{\mathbf{s}_\infty}) \text{ (notation of 1.7, 1.8).}$$

1.12. Let $X = Z_{J, y, \delta}$ or $\bar{G}$. There is a well-defined map from the irreducible components of $\Lambda(X)$ to the nilpotent conjugacy classes of $\mathfrak{g}^*$ which sends the irreducible component C of $\Lambda(X)$ to the nilpotent conjugacy class $\mathcal{O}$, where $\mu_{G \times G}(C) \cap \{(f, -f) \in \mathcal{O} \times \mathcal{O}\}$ is dense in $\mu_{G \times G}(C)$.

REFERENCES

- [Gi] V. Ginzburg, *Admissible modules on a symmetric space*, Astérisque **173-174** (1989), 199-256.
- [H1] X. He, *The G -stable pieces of the wonderful compactification*, math.RT/0412302.
- [H2] X. He, *The character sheaves on the group compactification*, math.RT/0508068.

- [KS] M. Kashiwara and P. Schapira, *Sheaves on manifolds*, Springer-Verlag, Berlin, 1994.
- [L1] G. Lusztig, *Parabolic character sheaves I*, Moscow Math.J **4** (2004), 153-179.
- [L2] G. Lusztig, *Parabolic character sheaves II*, Moscow Math.J. **4** (2004), 869-896.
- [MV] I. Mirković and K. Vilonen, *Characteristic varieties of character sheaves*, Invent.Math. **93** (1988), 405-418.
- [S] T. A. Springer, *Intersection cohomology of $B \times B$ -orbit closures in group compactifications*, J.Alg. **258** (2002), 71-111.

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON, NJ 08540
E-mail address: hugo@math.ias.edu

DEPARTMENT OF MATHEMATICS, M.I.T., CAMBRIDGE, MA 02139
E-mail address: gyuri@math.mit.edu